



Neuropsychological functioning and social anhedonia: Three-year follow-up data from a longitudinal community high risk study

Alex S. Cohen^a, Shannon M. Couture^b, Jack J. Blanchard^{c,*}

^a Louisiana State University, Department of Psychology, United States

^b University of Southern California, Department of Psychology, United States

^c University of Maryland, College Park, Department of Psychology, MD 20742 4411, United States

ARTICLE INFO

Article history:

Received 23 August 2011

Received in revised form

12 March 2012

Accepted 22 March 2012

Keywords:

Social anhedonia

Schizophrenia

Schizotypy

Prodrome

Risk

Neurocognitive

Neuropsychological

ABSTRACT

Social anhedonia is a promising vulnerability marker for schizophrenia-spectrum pathology. Prior research has demonstrated that individuals with psychometrically-defined social anhedonia show a range of “schizophrenia-like” neurocognitive abnormalities. However, this research is limited in that it is based largely on the study of college students. The present article reports findings from a longitudinal study of social anhedonia recruited from a community sample. As part of this study, a neurocognitive battery was administered at baseline and at three-year follow-up sessions to participants with ($n = 78$) versus without ($n = 77$) social anhedonia. Additional measures of global functioning and schizotypal, schizoid and paranoid schizophrenia-spectrum symptoms were also administered. Across groups, subjects showed significant improvement in neurocognitive functioning over time. Compared to controls, at follow-up, individuals with social anhedonia showed significantly poorer attentional vigilance and simple processing speed, but failed to evidence impairments in immediate or delayed verbal memory, immediate or delayed visual memory, visual or verbal working memory, olfaction or executive abilities. At follow-up, within the social anhedonia group, schizoid (and to a lesser extent, schizotypal) symptom severity was associated with a range of neurocognitive impairments. Neurocognitive impairments were generally not associated with paranoid symptoms or global functioning. Baseline neurocognitive performance was not significantly predictive of follow-up symptom severity or functioning. Collectively, these findings suggest that neurocognitive dysfunctions only characterize a subset of individuals with social anhedonia.

© 2012 Elsevier Ltd. All rights reserved.

1. Introduction

Social anhedonia, defined in terms of an inability to experience pleasure from social interactions, may be a risk marker for schizophrenia-spectrum disorders. Social anhedonia appears to be modestly heritable (Cohen et al., 2010; Kendler et al., 1996), is associated with reduced functioning (Blanchard et al., 2011), and has been associated with the emergence of schizophrenia-spectrum disorders in longitudinal questionnaire-based psychometric “high risk” paradigms using college student participants (e.g., Gooding et al., 2005; Kwapił, 1998). Cross-sectional studies have also found that socially anhedonic individuals evidence a number of “schizophrenia-like” neurocognitive abnormalities, albeit in attenuated form. For example, individuals with elevated levels of social anhedonia have demonstrated impairments in

visual-spatial working memory (Gooding and Tallent, 2003, 2004; Tallent and Gooding, 1999), visual-spatial construction (Gooding and Braun, 2004), visual-spatial delayed memory (Gooding and Tallent, 2004), sustained visual attention (Gooding et al., 2006) and executive functioning (Tallent and Gooding, 1999) compared to non-anhedonic controls. Given the importance of neurocognitive abnormalities for understanding schizophrenia (Green, 1996) and schizotypy (e.g., Gooding and Tallent, 2003, 2004; Tallent and Gooding, 1999) more generally, and the potential use of neurocognitive measures as an indicator of schizophrenia vulnerability (Gur et al., 2007), there is considerable merit to clarifying the relationship between neurocognitive impairments and social anhedonia.

A significant limitation of much prior research on neurocognitive functioning in individuals with social anhedonia is the reliance on non-representative college samples. Consider that these samples are typically composed of Caucasian college students from universities with above average intelligence (e.g., Gooding and Tallent, 2003). The study of samples that are college-educated may limit our

* Corresponding author. Tel.: +1 301 405 8438; fax: +1 301 405 0367.

E-mail addresses: jblancha@umd.edu, acohen@lsu.edu (J.J. Blanchard).

understanding of cognitive deficits in social anhedonia. The Maryland Longitudinal Study of Schizotypy (MLSS; Blanchard et al., 2011) was designed to address these concerns by applying the psychometric high risk paradigm to a representative community sample. In a prior article, we examined baseline differences in neurocognitive functioning across a wide array of attentional, memory, visual-spatial and language domains between individuals with social anhedonia and controls (Cohen et al., 2006). Our findings suggested that individuals with social anhedonia performed significantly worse than controls on tests of visual-spatial working memory, visual delayed memory and visual-constructional processing, but not on tests of verbal working memory, immediate or delayed memory, attentional vigilance or vocabulary ability. Moreover, neurocognitive performance was not significantly associated with functioning or severity of schizoid, schizotypal or paranoid symptoms, suggesting that neurocognitive abnormalities had not manifested in poorer clinical presentation at baseline. Given that these subjects were each 18-years old when they were recruited, and thus just entering the window of risk for onset of schizophrenia-spectrum disorders, it is important to understand how these neurocognitive impairments potentially changed over time.

The present study reports data from a three-year follow-up assessment from the MLSS. At the follow-up assessment, we repeated administration of the neurocognitive measures from the baseline assessment. This allowed us to determine the extent to which performance changed as individuals progressed through the risk period for schizophrenia-spectrum pathology. The longitudinal design also allowed us to examine whether baseline individual differences in neurocognitive performance were predictive of clinical outcomes at follow-up. We also expanded the scope of neurocognitive measures to include those tapping processing speed, executive functions and olfaction identification abilities – abilities not assessed at baseline but thought to be associated with schizophrenia vulnerability (e.g., Gooding et al., 1999). Olfaction identification is a particularly important neurocognitive ability to examine given recent claims that it reflects a valid vulnerability marker of schizophrenia-spectrum pathology (Turetsky et al., 2009). Finally, we administered measures of functioning and schizophrenia-spectrum symptoms at the follow-up assessment, which allowed us to evaluate whether neurocognitive impairment was related to functioning and schizophrenia-spectrum symptoms at the three-year follow-up assessment. For more detailed information on diagnosis, symptom severity and general functioning at follow-up in the social anhedonia group, the reader is referred to a companion article (Blanchard et al., 2011).

2. Methods

2.1. Participants

During the baseline screening, a cohort of 18-year old individuals ($N = 3508$) who lived within a 20-mile radius of the University of Maryland, College Park campus was identified using random-digit-dial methods. These individuals were each mailed a consent form and a screening self-report measure that included items from the Chapman schizotypy scales: the Revised-Social Anhedonia (RSAS; Eckblad et al., 1982), Perceptual Aberrations (PerAb; Chapman et al., 1978), Magical Ideation (MagId; Eckblad and Chapman, 1983) and Infrequency (Chapman and Chapman, 1976) scales. Response rate was high ($n = 2434$; 69%). Extreme scorers on the RSAS, selected as potential candidates for the social anhedonia group, were defined as either having a) an RSAS score 1.9 standard deviations above their respective gender and ethnicity-centered means (Chapman et al., 1994; Gooding et al., 2005; Kwapil, 1998), and/or b) having a Bayesian probabilities of belonging to the social

anhedonia taxon greater or equal to .50 using Maximum Covariate Analysis taxometric method (MAXCOV; See Horan et al., 2004 for a review of this methodology). A cutoff score of 1.9 identifies less than 3% of the population, which is conservative based on the purported 10% prevalence of social anhedonia (e.g., Horan et al., 2004). Using these methods, 86 socially anhedonic individuals were identified who agreed to participate at baseline. Additionally, 89 controls were identified based on their scores on the RSAS, Perceptual Aberrations, and Magical Ideation scales being below .50 standard deviations from the gender and ethnicity derived group means, and Bayesian probabilities of taxon membership less than .50 using the MAXCOV taxometric procedure. There were no statistically significant differences in neurocognitive functioning between subjects identified using extreme scores versus taxon membership. The control participant group was matched to the social anhedonia group on gender and race variables. The follow-up assessment was conducted approximately three years after the completion of the baseline assessment. Retention from the baseline study was similarly excellent between the social anhedonia (91%) and control (87%) groups. For this study, neurocognitive data were available for 78 social anhedonic participants and 77 controls. The descriptive data for these groups are presented in Table 1.

2.2. Psychometric schizotypy measures

The RSAS (Eckblad et al., 1982), was used to measure social anhedonia. The RSAS is a 40 item true–false self-report questionnaire designed to measure deficits in social pleasure. The PerAb (Chapman et al., 1978) and the MagId (Eckblad and Chapman, 1983) scales were used to measure psychosis proneness to screen controls. The PerAb is a 35 item true–false self-report questionnaire designed to measure distortions in the perception of one's own body and environment. The MagId is a 30 item true–false self-report questionnaire designed to measure beliefs about causation that deviate from the norm. The RSAS, PerAb and MagId each have documented validity and reliability, and the reader is referred to their source documents, referenced above, for their psychometric properties.

Table 1

Follow-up demographic and clinical data for social anhedonics ($n = 78$) and controls ($n = 77$).^a

	Social anhedonia	Controls
Gender		
% Female	45 (58%)	40 (52%)
Ethnicity		
Caucasian	35 (45%)	37 (48%)
African-American	37 (47%)	30 (39%)
Hispanic	4 (5%)	7 (9%)
Asian	1 (1%)	2 (3%)
Other/refused	1 (1%)	1 (1%)
Age	21.45 ± .50	21.52 ± .53
Epoch from baseline to follow-up (in days)	1021.37 ± 74.84	1082.78 ± 46.41
IPDE schizophrenia-spectrum scores^b		
Schizotypal	.53 ± 1.00	.08 ± .48
Schizoid	.77 ± 1.18	.10 ± .53
Paranoid	.65 ± 1.04	.12 ± .54
Overall functioning		
Revised Social Anhedonia Scale ^b	15.13 ± 6.99	6.58 ± 3.93
Global Assessment of Functioning ^b	73.76 ± 13.62	83.30 ± 10.73
Treatment history:^b		
Mood disorder diagnosis	34 (44%)	16 (21%)
Substance abuse diagnosis	18 (23%)	21 (27%)
Psychosis diagnosis	1 (1%)	1 (1%)
Inpatient treatment ^b	7 (9%)	1 (1.3%)
Outpatient treatment ^b	28 (36%)	15 (20%)

^a Three participants who met criteria for a lifetime psychotic disorder at baseline were excluded.

^b Data from the follow-up assessment.

The Infrequency Scale (Chapman and Chapman, 1976) was included to determine the extent to which participants' responses were valid. The infrequency scale is a 17 item true–false questionnaire. As in other studies (e.g., Luh and Gooding, 1999), individuals who endorsed three or more infrequency items were excluded from participation in this study.

2.3. Diagnostic interviews

Participants were administered a series of semi-structured diagnostic interviews during their baseline and follow-up assessments. The mood, psychosis, and substance abuse modules from the Structured Clinical Interview for the Diagnostic and Statistical Manual for Mental Disorders, Fourth edition (American Psychiatric Association, 1994) were administered to evaluate Axis I diagnoses. Schizophrenia-spectrum symptomatology was determined using the schizoid, schizotypal and paranoid personality disorder modules from the International Personality Disorders Examination (Loranger et al., 1994). Dimensional scores, computed as a sum of the respective ratings from the schizoid, schizotypal and paranoid modules, were computed and examined in this study. Diagnostic interviews were administered by doctoral-level graduate students who had received rigorous training in both SCID and IPDE administration. Supervision was provided by two doctoral psychologists (J. Blanchard & A. Cohen) with extensive experience in SCID and IPDE administration. Final diagnoses and ratings were discussed during monthly case conferences until consensus diagnoses had been reached between the case conference group members, which included the principle investigator (J. Blanchard) and at least two doctoral-level graduate students who reviewed the videotaped interviews. The participants, interviewers and consensus judges were each blind to the participants' group classification. As in our baseline study (Cohen et al., 2006), one participant from the social anhedonia and two from the control groups met criteria for psychotic disorders and were excluded from the studies. Individuals who developed psychotic symptoms after baseline were included in the present study. The rationale for this was that we were primarily interested in individuals at elevated risk for schizophrenia as opposed to those already meeting criteria for the disorder. The descriptive data for lifetime diagnoses are presented in Table 1.

2.4. Neuropsychological measures

Measures from the baseline assessment, selected based on prior investigations of neuropsychological functioning in schizophrenia and individuals at-risk for schizophrenia more generally (Heinrichs and Zakzanis, 1998), were readministered during the follow-up session. Memory tests included: the total recall scores from the Logical Memory I and II tests (Wechsler, 1987, 1997) as measures of immediate and delayed verbal memory respectively and the total correct scores from the Visual Reproduction I and II tests (Wechsler, 1987) as measures of immediate and delayed visual-spatial memory respectively. Working memory was assessed with the following tests: the total score from the Digit Span test (Wechsler, 1987) as a measure of immediate memory ability, the total correct score from the Spatial Span test (Wechsler, 1987) as a measure of visual-spatial attention/working memory, and the total correct from the Letter Number Sequencing test (Wechsler, 1987) as a measure of verbal working memory. The Degraded Stimuli – Continuous Performance Test (DSCPT; Nuechterlein and Asarnow, 1992), a measure of sustained attention was also used. Raw scores, as opposed to age corrected scores, were used for each of the Wechsler variables because all of the participants in this study were in the same age range, and because

raw scores provide greater variability. Increasing scores for each of these variables reflects better performance. The block design and vocabulary tests (Wechsler, 1997), which were administered as proxies for general cognitive ability at baseline, were not re-administered at follow-up.

We introduced three additional measures for the follow-up assessment. First, to expand the measures of executive functions, we included the total correct and number of perseverative errors variables from the Wisconsin Card Sorting Task (WCST; Heaton & PAR Staff, 1999) and the total seconds to completion from the Trails B test (a.k.a. “complex” processing speed; Boll, 1981). As a measure of simple processing speed, we included total seconds to completion from the Trails A (Boll, 1981). Finally, we administered a measure of olfaction recognition (UPSIT; Doty, 1984). Missing data included: one control and one social anhedonic subject for the WCST, one control and two social anhedonics for the UPSIT, and one control and four social anhedonics for the DSCPT.

2.5. General functioning

Participants' general level of functioning was measured using the Global Assessment of Functioning (GAF; American Psychiatric Association, 1994) scale. GAF scores, based on a scale from one to 100, reflect a measure of an individual's psychological, social and occupational functioning for the prior month. Increasing scores reflect better functioning. The team consensus approach, outlined above, was used to assign GAF scores for each case.

2.6. Socio-economic status

Socio-economic status for the participants' father and mother was separately determined using the Hollingshead scale (Hollingshead, 1975). When both scores were available, the two scores were averaged together.

2.7. Statistical analyses

Analyses were conducted in five steps. First, we compared the social anhedonia and control groups in descriptive characteristics to ensure there were no confounding variables worthy of consideration in the subsequent analyses. Second, we employed a series of time (i.e., baseline and follow-up) by group (i.e., social anhedonia versus controls) repeated-measure ANOVAs to evaluate neurocognitive performance over time across the two groups. Third, we compared the social anhedonia and control groups on performance for the measures that were only administered at follow-up, specifically those tapping executive, processing speed and olfaction abilities, using a series of *t*-tests. Fourth, in order to determine the extent to which neurocognitive performance was associated with symptom and functioning variables within the social anhedonia group, bivariate correlations were computed between the neurocognitive test scores and the paranoid, schizoid and schizotypal and GAF scores for the social anhedonia group only. Finally, we examined the relationship between baseline neurocognitive measures and follow-up schizotypal symptom and functioning scores. This analysis helped evaluate the predictive power of neurocognitive performance in terms of symptom severity and functioning. The paranoid, schizoid and schizotypal dimensional scores from the IPDE were square root transformed to compensate for excessive positive skew (skew > 1.5). Unless otherwise noted, all variables are normally distributed and all statistical tests are two-tailed.

3. Results

3.1. Descriptive and clinical characteristics

Descriptive statistics were computed and compared between the social anhedonia and control groups for follow-up data (see Table 1). The groups were not statistically different in sex, ethnicity, age, time between baseline and follow-up assessments, or history of substance abuse or schizophrenia diagnoses (p 's > .10). The social anhedonia group had significantly greater schizotypal (t [153] = 3.54, p = .001), schizoid (t [153] = 4.51, p < .001) and paranoid (t [153] = 4.02, p < .001) ratings, poorer functioning (t [153] = 4.84, p < .001), higher frequency of inpatient (χ^2 [1] = 4.66, p = .03) and outpatient (χ^2 [1] = 5.21, p = .02) treatment history and history of major depressive disorder (χ^2 [1] = 9.23, p = .002). For more information regarding these variables, the author is referred to Blanchard et al. (2011).

3.2. Group differences in neurocognitive performance: baseline to follow-up

Table 2 contains the means, standard deviations, and test statistics for the social anhedonia and control groups for neurocognitive performance across baseline and follow-up sessions. Significant time effects indicated that, across groups, performance significantly improved across the three-year epoch in immediate and delayed verbal and visual memory, and for verbal working memory. Significant main effects for group, indicated that, compared to controls, the social anhedonia group was significantly poorer in delayed visual memory and attentional vigilance performance, but the groups did not differ on any of the other tasks. There were no significant group by time interactions. Our expectation that individuals with social anhedonia would show poorer neurocognitive functioning was generally not supported.

It is noteworthy that we reported baseline group differences in visual memory and attentional vigilance performance in our prior publication of baseline differences (Cohen et al., 2006), yet no baseline differences were observed in the present analyses of baseline performance (which only contains those individuals who completed the follow-up). This disparity is because the five individuals with social anhedonia lost to the follow-up had significantly poorer neurocognitive functioning on these tasks than those

Table 3
Neurocognitive measures only available at follow-up.

	Control		Social anhedonia		t	d
	M	SD	M	SD		
WCST (total correct)	68.92	9.32	69.51	8.57	.41	.07
WCST (perseverative errors)	9.22	10.73	10.50	8.65	.82	.13
Olfaction functioning	33.26	4.05	34.66	3.37	2.35*	.38
Simple processing speed (Trails A)	25.14	9.63	28.85	12.14	2.13*	.34
Complex processing speed (Trails B)	56.52	19.14	62.96	30.07	1.61	.26

* = p < .05.

retained in the study (t 's[83] = 2.44 and 2.96, p 's = .02 and .004, respectively).

3.3. Group differences in neurocognitive task performance assessed at follow-up

Table 3 contains means, standard deviations and test statistics for those tests administered only at follow-up for the social anhedonia and control groups. The social anhedonia group was significantly slower on the processing speed task and significantly more accurate on the olfaction recognition test. In general, expected group differences were not seen with the additional neurocognitive tasks assessed at follow-up.

3.4. The relationship between global functioning, symptoms and neurocognitive performance (within the social anhedonia group) at follow-up

Correlations computed between the functioning, schizotypal, schizoid and paranoid symptom and neurocognitive performance scores are in Table 4. Increasing schizoid symptoms corresponded to poorer immediate and delayed verbal and visual memory, poorer olfaction recognition and slower processing speed for both simple (i.e., Trails A) and complex (i.e., Trails B) processing speed tasks. Severity of schizotypal symptoms was associated with poorer immediate verbal memory and olfaction recognition. Neither functioning nor paranoid symptoms were significantly associated with any neurocognitive performance variables. In sum, schizoid and, to a lesser extent, schizotypal, but not paranoid symptoms were associated with a range of neurocognitive impairments.

Table 2
Neurocognitive performance between baseline and three-year follow-up administrations for controls versus social anhedonia (S. A.) groups.

	Time	Control	Social anhedonia	F			d		
		$M \pm SD$	$M \pm SD$	Time	Between	Interaction	Time	Between: control	Between: S. A.
Immediate verbal memory	Baseline	44.31 ± 9.89	42.63 ± 10.29	21.51***	1.06	.04	.17	-.35	-.35
	Follow-up	47.84 ± 10.17	46.46 ± 11.60				.13		
Delayed verbal memory	Baseline	27.88 ± 7.59	27.14 ± 7.24	51.81***	1.29	1.08	.10	-.53	-.40
	Follow-up	31.84 ± 7.41	30.10 ± 7.49				.23		
Immediate visual memory	Baseline	93.68 ± 7.95	91.56 ± 10.90	27.35***	1.89	.55	.22	-.43	-.49
	Follow-up	96.97 ± 7.38	95.95 ± 6.96				.14		
Delayed visual memory	Baseline	84.70 ± 13.90	80.54 ± 16.06	20.67***	4.63*	.01	.28	-.41	-.36
	Follow-up	90.29 ± 13.35	85.94 ± 14.24				.32		
Verbal working memory (LN Seq)	Baseline	11.66 ± 2.74	11.05 ± 2.56	3.24	1.27	1.02	.23	-.05	-.18
	Follow-up	11.79 ± 2.73	11.51 ± 2.63				.10		
Visual working memory (spatial span)	Baseline	17.52 ± 3.30	17.31 ± 3.12	.27	1.38	2.94	.07	-.08	.14
	Follow-up	17.77 ± 3.31	16.85 ± 3.33				.28		
Verbal working memory (digit span)	Baseline	18.30 ± 4.24	17.45 ± 3.88	12.54***	2.22	.07	.21	-.21	-.19
	Follow-up	19.19 ± 4.16	18.22 ± 4.06				.24		
Attentional vigilance: d prime	Baseline	2.30 ± .91	2.03 ± .95	.37	4.77*	.91	.26	-.02	.13
	Follow-up	2.32 ± .95	1.93 ± 1.09				.38		

* = p < .05, *** = p < .001.

Table 4
Bivariate correlations between clinical and neurocognitive functioning variables at follow-up, within the social anhedonia group.

	Functioning	IPDE schizophrenia-spectrum traits		
	GAF	Schizotypy	Schizoid	Paranoia
Immediate verbal memory	.22	-.24*	-.33**	-.14
Delayed verbal memory	.12	-.18	-.29*	-.10
Immediate visual memory	-.04	.01	-.25*	-.01
Delayed visual memory	.02	-.02	-.22*	.00
Verbal working memory (LN Seq)	-.17	-.12	.00	.11
Visual working memory (spatial span)	.16	-.19	-.19	-.21
Verbal working memory (digit span)	-.01	-.11	-.11	.04
WCST (total correct)	-.08	-.02	.03	-.03
WCST (perseverative errors)	.04	.00	-.09	.02
Olfaction functioning	.11	-.24*	-.25*	.08
Simple processing speed (Trails A)	-.08	.18	.25*	.08
Complex processing speed (Trails B)	-.11	.18	.24*	-.02
Attentional vigilance: <i>d</i> prime	-.04	-.02	-.18	-.04

Note – Schizophrenia-spectrum scores square root transformed to compensate for excessive skew.

* = $p < .05$, ** = $p < .01$.

Neurocognitive variables were not significantly related to functioning as hypothesized.

3.5. The relationship between baseline neurocognition and follow-up symptom and functioning measures

Correlations computed between the baseline neurocognitive measures and the follow-up schizotypal symptoms and GAF scores were then computed. Poorer visual-spatial working memory performance was associated with more severe follow-up paranoia symptoms ($r[78] = -.28$, $p = .01$). Beyond this, baseline neurocognition performance was not significantly associated with any of the schizotypal, paranoid or schizoid symptom scores (p 's $> .05$). With respect to functioning, none of the baseline neurocognition scores were significantly associated with any of the follow-up functioning scores in the expected direction. In sum, baseline neurocognition held little power in predicting follow-up schizotypal symptoms or functioning.

4. Discussion

This study examined neurocognitive functioning in individuals with psychometrically-defined social anhedonia and controls. There are four relatively unique features of this study. First, in contrast to most studies of this kind that recruit from college student populations, participants were recruited from a large metropolitan region. Second, due to the longitudinal design of this study and the excellent retention for study participants, we were able to examine potential changes in neurocognitive performance across a three-year epoch. Third, we employed a relatively large battery of standard neurocognitive instruments tapping a broad range of abilities. Finally, we employed measures of functioning and schizophrenia-spectrum pathology, thus offering insight into the correlates of neurocognitive impairments in individuals with social anhedonia.

In contrast to our expectations, there were very few neurocognitive impairments in the social anhedonia group at follow-up. Only two measures – one tapping attentional vigilance and the other simple processing speed, were significantly different between the groups at follow-up. In contrast, immediate and delayed memory across both verbal and visual domains, working memory, olfaction recognition and a range of executive functions were preserved in the social anhedonia group. These findings are inconsistent with what was observed at baseline, where the social anhedonia group demonstrated a broader range of visual memory

impairments. The reasons for this seeming discrepancy reflect that the five subjects lost to follow-up tended to have poorer visual memory, and that the group differences (which were at a small effect size at baseline) narrowed over time. Our findings are inconsistent with many published studies on neurocognition in college students with social anhedonia (e.g., Gooding and Tallent, 2004), and the reasons for this disparity are unclear at the present time. It is worth noting that there are a number of null findings with respect to neurocognition in psychometrically-defined schizotypy more generally (e.g., Lenzenweger and Gold, 2000; Cohen et al., 2009) – so the present findings are not wholly inconsistent with the extant literature. For what it's worth, it is clear that the samples employed in the present study were much more representative of the general US population in terms of ethnicity, education, substance use/abuse and global cognitive ability (see Blanchard et al., 2011 and Cohen et al., 2006), though it is unclear if sample differences between this and prior social anhedonia studies may have contributed to the null findings. Regardless, care is recommended in the assumption of generalizability from college samples to more representative community populations.

The present data did not support the notion that neurocognitive performance was, for the most part, predictive of functioning. Interestingly, this finding is consistent with that reported by Gooding et al. (2005) in that baseline laboratory metrics were not predictive of symptom severity after a five year epoch. In that study, plexus visibility, smooth pursuit eye tracking and antisaccade task errors were the metrics of interest. However, our findings are inconsistent with what is seen in schizophrenia, as neurocognitive dysfunction has been shown to be predictive of functioning and illness severity over time (e.g., Milev et al., 2005). Clarifying why neurocognition and illness course are related in patients with schizophrenia, but not in individuals with social anhedonia is a potentially interesting future research line.

It is noteworthy that neurocognitive performance was relatively poor in some socially anhedonic individuals. At follow-up, within the social anhedonia group, severity of schizoid symptoms was associated with poorer neurocognitive performance in a range of domains, including immediate and delayed verbal and visual memory, olfaction recognition and processing speed. Moreover, severity of schizotypal symptoms was associated with poorer performance on immediate verbal memory and olfaction recognition tests. This suggests that a subset of individuals with social anhedonia – those that are manifesting schizoid and schizotypal symptoms, show neurocognitive dysfunctions. It is curious that

schizoid symptoms, as opposed to paranoid symptoms were associated with poorer neurocognitive performance. While the meaning of this finding is unclear at the present time, it echoes findings from the schizophrenia literature that patients with prominent negative symptoms have poorer neurocognitive functioning than patients with positive or disorganization symptoms (Cohen et al., 2007). Thus, schizoid symptoms may reflect a marker of neurocognitive pathology in individuals with social anhedonia. The nature of the relationship between schizotypal symptoms and neurocognitive deficits is unclear at the present time. Further examining the relationship between these variables, and whether they share pathophysiological substrata, reflects an important line for future research.

Methodological issues may have contributed to two of the major null findings in this study. First, in contrast to what is seen in the literature on schizophrenia (Green, 1996), global functioning was not significantly related to neurocognitive performance. This is an important finding because it suggests that, in individuals with social anhedonia, global functioning impairments do not manifest as a function of neurocognitive impairments. However, this null finding might reflect the measure of functioning employed not being particularly sensitive and reflecting an overly broad range of symptom, occupational and social domains. For this reason, examining the functional sequelae of social anhedonia as they independently manifest across a broad range of social, academic and occupational domains is potentially important for future research. Second, although the neurocognitive tests employed in this study are commonly used in clinical practice and schizophrenia research more generally, they may be insufficiently sensitive for detecting differences in at-risk individuals. It seems unlikely that insensitivity wholly accounts for the lack of group differences in neurocognitive functioning, as there was sufficient variability in performance across individuals with social anhedonia to be significantly correlated to schizoid and schizotypal symptoms. Nonetheless, further research, perhaps employing methods derived from cognitive sciences (MacDonald and Carter, 2002), would address this issue. Finally, while this study had excellent subject retention from the baseline to follow-up assessments, a handful of subjects from the social anhedonia group were lost due to attrition. It appeared to be the case that, at least in some respects, these individuals had relatively poor neurocognitive functioning. Moreover, it stands to reason that they had a higher risk of manifesting schizophrenia.

In summary, results from the three-year follow-up assessment of neurocognitive functioning from subjects in the MLSS provide little evidence of impairments in individuals with social anhedonia. It is noteworthy that subjects in this study were young adults and still progressing through the “window of risk”. Further longitudinal assessment of these individuals will hopefully provide further insights into the neurocognition of social anhedonia.

Role of the funding source

Funding for this study was provided by NIMH Grants R01MH51240 and K02MH079231 to Jack Blanchard; the NIMH had no further role in study design; in the collection, analysis and interpretation of data; in the writing of the report; and in the decision to submit the paper for publication.

Contributors

All three authors were involved in the literature search, statistical analysis, interpretation of the results and writing of the manuscript. Jack Blanchard was the principal investigator of the project and designed the methodology.

Conflicts of interest

There are no conflicts of interest to report.

Acknowledgments

The authors wish to acknowledge the efforts of Monica Mann, Courtney Forbes and Jay Carreno, who helped collect data for this project. We also acknowledge the efforts of the research participants who made this project possible.

References

- American Psychiatric Association. Diagnostic and Statistical Manual of mental disorders (DSM-IV). Washington, DC: American Psychiatric Press; 1994.
- Blanchard JJ, Collins LM, Aghevli M, Leung WW, Cohen AS. Social anhedonia and schizotypy in a community sample: the Maryland longitudinal study of schizotypy. *Schizophrenia Bulletin* 2011;37:587–602.
- Boll TJ. The Halstead-Reitan neuropsychological battery. In: Filskov SB, Boll TJ, editors. *Handbook of clinical neuropsychology*. New York: Wiley-Interscience; 1981. p. 577–607.
- Chapman LJ, Chapman JP, Kwapil TR, Eckblad M, Zinser MC. Putatively psychosis-prone subjects 10 years later. *Journal of Abnormal Psychology* 1994;103:171–83.
- Chapman LJ, Chapman JP, Raulin Misholve L. Perceptual Aberration scale. University of Wisconsin—Madison: Unpublished manuscript; 1978.
- Chapman LJ, Chapman JP. The infrequency scale. p. Unpublished Test. 1976.
- Cohen AS, Emmerson LC, Mann MC, Forbes CB, Blanchard JJ. Schizotypal, schizoid and paranoid characteristics in the biological parents of social anhedonics. *Psychiatry Research* 2010;178:79–83.
- Cohen AS, Leung WW, Saperstein AM, Blanchard JJ. Neuropsychological functioning and social anhedonia: results from a community high-risk study. *Schizophrenia Research* 2006;85:132–41.
- Cohen AS, Saperstein AM, Gold JM, Kirkpatrick B, Carpenter Jr WT, Buchanan RW. Neuropsychology of the deficit syndrome: new data and meta-analysis of findings to date. *Schizophrenia Bulletin* 2007;33:1201–12.
- Cohen AS, Iglesias B, Minor KS. The neurocognitive underpinnings of diminished expressivity in schizotypy: what the voice reveals. *Schizophrenia Research* 2009;109:38–45.
- Doty RL. Development of the University of Pennsylvania smell identification test: a standardized microencapsulated test of olfactory function. *Physiological Behavior* 1984;32:489–502.
- Eckblad M, Chapman LJ. Magical ideation as an indicator of schizotypy. *Journal of Consulting and Clinical Psychology* 1983;51:215–25.
- Eckblad ML, Chapman LJ, Chapman JP, Mishlove M. The revised social anhedonia scale. Madison: University of Wisconsin; 1982.
- Gooding DC, Braun JG. Visuoconstructive performance, implicit hemispatial inattention, and schizotypy. *Schizophrenia Research* 2004;68:261–9.
- Gooding DC, Kwapil TR, Tallent KA. Wisconsin card sorting test deficits in schizotypic individuals. *Schizophrenia Research* 1999;40:201–9.
- Gooding DC, Matts CW, Rollmann EA. Sustained attention deficits in relation to psychometrically identified schizotypy: evaluating a potential endophenotypic marker. *Schizophrenia Research* 2006;82:27–37.
- Gooding DC, Tallent KA. Spatial, object, and affective working memory in social anhedonia: an exploratory study. *Schizophrenia Research* 2003;63:247–60.
- Gooding DC, Tallent KA. Nonverbal working memory deficits in schizophrenia patients: evidence of a supramodal executive processing deficit. *Schizophrenia Research* 2004;68:189–201.
- Gooding DC, Tallent KA, Matts CW. Clinical status of at-risk individuals 5 years later: further validation of the psychometric high-risk strategy. *Journal of Abnormal Psychology* 2005;114:170–5.
- Green MF. What are the functional consequences of neurocognitive deficits in schizophrenia? *American Journal of Psychiatry* 1996;153:321–30.
- Gur RE, Calkins ME, Gur RC, Horan WP, Nuechterlein KH, Seidman LJ, et al. The consortium on the genetics of schizophrenia: neurocognitive endophenotypes. *Schizophrenia Bulletin* 2007;33:49–68.
- Heaton RK, PAR Staff. Wisconsin card sorting test: computer version 3 for Windows. Odessa, FL: Psychological Assessment Resources (PAR); 1999.
- Heinrichs RW, Zakzanis KK. Neurocognitive deficit in schizophrenia: a quantitative review of the evidence. *Neuropsychology* 1998;12:426–45.
- Hollingshead AB. Four-factor index of social status. New Haven, CT: Yale Station; 1975.
- Horan WP, Blanchard JJ, Gangestad SW, Kwapil TR. The psychometric detection of schizotypy: do putative schizotypy indicators identify the same latent class? *Journal of Abnormal Psychology* 2004;113:339–57.
- Kendler KS, Thacker L, Walsh D. Self-report measures of schizotypy as indices of familial vulnerability to schizophrenia. *Schizophrenia Bulletin* 1996;22:511–20.
- Kwapil TR. Social anhedonia as a predictor of the development of schizophrenia-spectrum disorders. *Journal of Abnormal Psychology* 1998;107:558–65.
- Lenzenweger MF, Gold JM. Auditory working memory and verbal recall memory in schizotypy. *Schizophrenia Research* 2000;42:101–10.
- Loranger AW, Sartorius N, Andreoli A, Berger P, Buchheim P, Channabasavanna SM, et al. The international personality disorder examination. The world health

- organization/alcohol, drug abuse, and mental health administration international pilot study of personality disorders. *Archives of General Psychiatry* 1994; 51:215–24.
- Luh KE, Gooding DC. Perceptual biases in psychosis-prone individuals. *Journal of Abnormal Psychology* 1999;108:283–9.
- MacDonald III AW, Carter CS. Cognitive experimental approaches to investigating impaired cognition in schizophrenia: a paradigm shift. *Journal of Clinical and Experimental Neuropsychology* 2002;24:873–82.
- Milev P, Ho B-C, Arndt S, Andreasen NC. Predictive values of neurocognition and negative symptoms on functional outcome in schizophrenia: a longitudinal first-episode study with 7-year follow-up. *American Journal of Psychiatry* 2005; 162:495.
- Nuechterlein KH, Asarnow RF. Manual and computer program for the UCLA continuous performance test: version 4; 1992.
- Tallent KA, Gooding DC. Working memory and Wisconsin card sorting test performance in schizotypic individuals: a replication and extension. *Psychiatry Research* 1999;89:161–70.
- Turetsky BI, Hahn CG, Borgmann-Winter K, Moberg PJ. Scents and nonsense: olfactory dysfunction in schizophrenia. *Schizophrenia Bulletin* 2009;35: 1117–31.
- Wechsler D. Wechsler memory scale-revised. New York, NY: Psychological Corporation; 1987.
- Wechsler D. The Wechsler adult intelligence scale – third edition. San Antonio, TX: The Psychological Corporation; 1997.